VOL. 433 JOURNAL OF CHROMATOGRAPHY DECEMBER 9, 1988

(Biomedical Applications, Vol. 77)

CONTENTS

(Abstracts/Contents Lists published in Analytical Abstracts, ASCA, Biochemical Abstracts, Biological Abstracts, Chemical Abstracts, Chemical Titles, Chromatography Abstracts, Clinical Chemistry Lookout, Current Contents/Physical, Chemical & Earth Sciences, Current Contents/Life Sciences, Deep-Sea Research/Part B: Oceanographic Literature Review, Excerpta Medica, Index Medicus, Mass Spectrometry Bulletin, PASCAL-CNRS, Pharmaceutical Abstracts, Referativnyi Zhurnal, Science Citation Index and Trends in Biotechnology)

Measurement of plasma oxalate by capillary gas chromatography and its validation by isotope dilution mass spectrometry
by N.C. France, P.T. Holland, T.K. McGhie and M.R. Wallace (Hamilton, New Zealand) (Received August 8th, 1988) 1

Screening, confirmation and quantification of boldenone sulfate in equine urine after administration of boldenone undecylenate (Equipoise™)
by L.O.G. Weidolf, T.M.P. Chichila and J.D. Henion (Ithaca, NY, U.S.A.) (Received August 8th, 1988) 9

Effects of long-term storage on the concentrations of the unconjugated acidic metabolites of the trace amines, indoleamines and catecholamines
by B.A. Davis (Saskatoon, Canada) (Received August 8th, 1988) 23

Selected-ion monitoring gas chromatographic–mass spectrometric analysis of catecholamines: enhancement of sensitivity by a simple clean-up step on Sephadex G-10
by G.K.E. Scriba and R.T. Borchardt (Lawrence, KS, U.S.A.) and J.A. Zirrolli and P.V. Fennessey (Denver, CO, U.S.A.) (Received August 5th, 1988) 31

Sensitive method for determination of picogram amounts of epinephrine and other catecholamines in microdissected samples of rat brain using liquid chromatography with electrochemical detection
by J. Opacka-Juffry, F. Tacconelli and C.W. Coen (London, U.K.) (Received July 7th, 1988) 41

Pre-column derivatization of amino acids with N,N-diethyl-2,4-dinitro-5-fluoroaniline and reversed-phase liquid chromatographic separation
by I. Fermo, F.M. Rubino, E. Bolzacchini, C. Arvelloni, R. Paroni and P.A. Bonini (Milan, Italy) (Received July 7th, 1988) 53

Assay of ATP-sulfurylase activity from rat liver by high-performance liquid chromatography
by M. Mina and E.F. Rossomando (Farmington, CT, U.S.A.) (Received August 16th, 1988) 63

(Continued overleaf)

Determination of erythrocyte porphyrins by reversed-phase high-performance liquid chromatography using capsule-type silica gels coated with silicone polymer
by T. Sakai, Y. Takeuchi, T. Araki and K. Ushio (Tokyo, Japan) (Received July 13th, 1988) 73

Quantitative evaluation of collagen fractions separated by acrylamide gel electrophoresis: its application for collagen typing in normal and pathological tissues
by G. Bellon and J.P. Borel (Reims, France) (Received July 27th, 1988) 81

Simultaneous determination of fentanyl and alfentanil in rat tissues by capillary column gas chromatography
by S. Björkman and D.R. Stanski (Stanford, CA, U.S.A.) (Received June 27th, 1988) 95

Enantioselective gas chromatographic assay with electron-capture detection for *dl*-fenfluramine and *dl*-norfenfluramine in plasma
by N.R. Srinivas, J.W. Hubbard, J.K. Cooper and K.K. Midha (Saskatoon, Canada) (Received August 22nd, 1988) 105

Determination of digoxin in serum by on-line immunoadsorptive clean-up high-performance liquid chromatographic separation and fluorescence-reaction detection
by E. Reh (Tutzing, F.R.G.) (Received July 28th, 1988) 119

Screening procedure for stimulant laxatives in urine using high-performance liquid chromatography with diode array detection
by R.O. Fullinfaw, R.W. Bury and R.F.W. Moulds (Parkville, Australia) (Received August 16th, 1988) 131

Determination of gossypol enantiomers in plasma after administration of racemate using high-performance liquid chromatography with precolumn chemical derivatisation
by D.-F. Wu, M.M. Reidenberg and D.E. Drayer (New York, NY, U.S.A.) (Received July 27th, 1988) 141

Liquid chromatographic and flow injection analysis of tetracycline using sensitized europium(III) luminescence detection
by T.J. Wenzel, L.M. Collette, D.T. Dahlen, S.M. Hendrickson and L.W. Yarmaloff (Lewiston, ME, U.S.A.) (Received July 19th, 1988) 149

High-performance liquid chromatographic determination of fusidic acid in plasma
by A. Rahman and N.E. Hoffman (Milwaukee, WI, U.S.A.) (Received August 23rd, 1988) 159

Determination of ambenonium in biological samples by reversed-phase ion-pair liquid chromatography
by K. Yamamoto, Y. Kohda, Y. Saitoh and F. Nakagawa (Tokyo, Japan) (Received July 20th, 1988) 167

High-performance liquid chromatographic determination of 6-amidino-2-naphthyl [4-(4,5-dihydro-1*H*-imidazol-2-yl)amino]benzoate dimethanesulphonate and its metabolites in biological fluids
by T. Marunaka, M. Maniwa, E. Matsushima, K.-I. Yoshida and R. Azuma (Tokushima, Japan) and M. Kurotori and S. Komatsu (Chiba, Japan) (Received August 2nd, 1988) 177

High-performance liquid chromatographic analysis of trimethoprim and sulfamethoxazole in microliter volumes of chinchilla middle ear effusion and serum
by G.R. Erdmann, D.M. Canafax and G.S. Giebink (Minneapolis, MN, U.S.A.) (Received August 1st, 1988) 187

Analysis of blood and urine samples for hydroxychloroquine and three major metabolites by high-performance liquid chromatography with fluorescence detection
by S.B. Williams, L.C. Patchen and F.C. Churchill (Atlanta, GA, U.S.A.) (Received July 27th, 1988) 197

High-performance liquid chromatographic analysis of desferrioxamine. Pharmacokinetic and metabolic studies
by T.P.A. Kruck, K. Teichert-Kuliszewska, E. Fisher, W. Kalow and D.R. McLachlan (Toronto, Canada) (Received July 15th, 1988) 207

Application of two-dimensional high-performance liquid chromatography in metabolism studies of ivermectin
by S.-H.L. Chiu, E. Sestokas and R. Taub (Rahway, NJ, U.S.A.) (Received July 19th, 1988) 217

Determination of tranexamic acid in human serum by high-performance liquid chromatography using selective pre-column derivatization with phenyl isothiocyanate
by K. Matsubayashi, C. Kojima and H. Tachizawa (Tokyo, Japan) (Received July 21st, 1988) 225

High-performance liquid chromatographic determination of a new nootropic, N-(2,6-dimethylphenyl)-2-(2-oxo-1-pyrrolidinyl)acetamide, in human serum and urine
by Y. Fujimaki, K. Sudo and H. Tachizawa (Tokyo, Japan) (Received June 10th, 1988) 235

Notes

Highly sensitive gas chromatographic–mass spectrometric method for the determination of platelet-activating factor in human blood
by K. Yamada, O. Asano, T. Yoshimura and K. Katayama (Ibaraki, Japan) (Received July 7th, 1988) 243

Improved procedure for preparation of pentafluorobenzyl derivatives of carboxylic acids for analysis by gas chromatography with electron-capture detection
by M.I. Daneshvar and J.B. Brooks (Atlanta, GA, U.S.A.) (Received August 22nd, 1988) 248

Fractionation of baboon chylomicrons and very-low-density lipoproteins by high-performance liquid chromatography
by M.C. Williams and R.S. Kushwaha (San Antonio, TX, U.S.A.) (Received July 12th, 1988) 257

Shielded hydrophobic phase: a new concept for direct injection analysis of biological fluids by high-performance liquid chromatography
by D.J. Gisch, B.T. Hunter and B. Feibush (Bellefonte, PA, U.S.A.) (Received July 5th, 1988) 264

Determination of stobadine, a novel cardioprotective drug, using capillary gas chromatography with nitrogen–phosphorus detection after its selective solid-phase extraction from serum
by V. Marko (Bratislava, Czechoslovakia) (Received July 21st, 1988) 269

(Continued overleaf)

Contents (continued)

Automated capillary gas chromatographic assay using flame ionization detection for the determination of topiramate in plasma
by M.L. Holland, J.A. Uetz and K.T. Ng (Spring House, PA, U.S.A.) (Received August 1st, 1988) 276

Determination of a new hypolipidemic agent, 2-(*trans*-4-isobutylcyclohexyl)-2-oxoethyl benzenesulphonate, and its metabolite by high-performance liquid chromatography and gas chromatography–mass fragmentography in human biological materials
by M. Maniwa, Y. Umeno and T. Marunaka (Tokuchima, Japan) (Received August 2nd, 1988) 282

Determination of 1,3-dimethyl-7-isobutylxanthine in guinea pig plasma by high-performance liquid chromatography
by O. Agostini, S. Manzini and M. Fedi (Florence, Italy) (Received July 13th, 1988) .. 288

Reversed-phase high-performance liquid chromatographic assay for the determination of the in vitro acetylation of *p*-aminobenzoic acid by human whole blood
by R.M. Lindsay (Dundee, U.K.), A.M. McLaren (Aberdeen, U.K.) and J.D. Baty (Dundee, U.K.) (Received July 6th, 1988) 292

Analysis of erythromycin and roxithromycin in plasma or serum by high-performance liquid chromatography using electrochemical detection
by N. Grgurinovich and A. Matthews (Adelaide, Australia) (Received July 8th, 1988) 298

Improved high-performance liquid chromatographic assay of antipyrine, hydroxymethylantipyrine, 4-hydroxyantipyrine and norantipyrine in urine
by M.A. Mikati, G.K. Szabo, R.J. Pylilo, B.W. LeDuc, T.R. Browne and D.J. Greenblatt (Boston, MA, U.S.A.) (Received August 18th, 1988) 305

Simultaneous determination of very low concentrations of haloperidol and reduced haloperidol in human serum by a liquid chromatographic method
by G.T. Vatassery, L.A. Herzan and M.W. Dysken (Minneapolis, MN, U.S.A.) (Received July 13th, 1988) 312

Reversed-phase high-performance liquid chromatographic assay to quantitate diastereomeric derivatives of metoprolol enantiomers in plasma
by D. Schuster, M. Woodruff Modi, D. Lalka and F.M. Gengo (Buffalo, NY, U.S.A.) (Received July 27th, 1988) 318

Rapid and sensitive high-performance liquid chromatographic assay for midazolam and 1-hydroxymidazolam, the major metabolite, in human serum
by A. Blackett, S. Dhillon, J.A. Cromarty and R. Horne (London, U.K.) and G. Richards (Southhall, U.K.) (Received July 25th, 1988) 326

Measurement of carprofen enantiomer concentrations in plasma and urine using L-leucinamide as the chiral coupling component
by H. Spahn (Frankfurt am Main, F.R.G. and San Francisco, CA, U.S.A.), I. Spahn, G. Plugmann and E. Mutschler (Frankfurt am Main, F.R.G.) and L.Z. Benet (San Francisco, CA, U.S.A.) (Received August 4th, 1988) 331

Determination of halofantrine and its principal metabolite desbutylhalofantrine in biological fluids by reversed-phase high-performance liquid chromatography
by K.A. Milton, S.A. Ward and G. Edwards (Liverpool, U.K.) (Received July 25th, 1988) 339

Simultaneous microassay for etretinate and its active metabolite, etretin, by reversed-phase high-performance liquid chromatography
by P. Thongnopnua and C.L. Zimmerman (Minneapolis, MN, U.S.A.) (Received July 11th, 1988) 345

Determination of tacrine hydrochloride in human serum by chloroform extraction, reversed-phase high-performance liquid chromatography and fluorimetric detection
by D.R. Forsyth, J.M. Ford, C.A. Truman, C.J.C. Roberts and G.K. Wilcock (Bristol, U.K.) (Received August 1st, 1988) 352

High-performance liquid chromatographic determination of nifurtimox in human serum
by C. Paulos, J. Paredes, I. Vasquez, G. Kunze and G. González-Martin (Santiago, Chile) (Received July 15th, 1988) 359

Chromatographic procedure for preparation of a high-purity C 1 esterase inhibitor concentrate
by A.G. Benny and C.M. Brimer (Private Bag, New Zealand) (Received August 8th, 1988) 363

Reversed-phase liquid chromatographic determination of chloroaluminum phthalocyanine tetrasulfonate in canine tissues and fluids
by M.M. Zuk and K. Tyczkowska (Raleigh, NC, U.S.A.), E. Ben-Hur (Beer-Sheva, Israel), H.C. Newman (Raleigh, NC, U.S.A.), I. Rosenthal (Bet-Dagan, Israel) and S.W. Crane (Raleigh, NC, U.S.A.) (Received August 23rd, 1988) 367

Letter to the Editor

Determination of phosphate, sulfate and oxalate in urine by ion chromatography
by R.P. Singh and G.H. Nancollas (Buffalo, NY, U.S.A.) (Received July 28th, 1988) .. 373

Related articles published in Journal of Chromatography, Vols. 454 and 455 377

Author Index 379

Subject Index 383

In articles with more than one author, the name of the author to whom correspondence should be addressed is indicated in the article heading by a 6-pointed asterisk (*)